

Appendix Table 1d. *Lifetime comorbidities of Generalized Anxiety Disorder with other anxiety, mood, and substance use disorders, in the NCS (United States)*

	Temporal Priority of GAD and Comorbid Disorder†								
	Lifetime Generalized Anxiety Disorder			Lifetime GAD First		Comorbid Disorder First		Both Started in Same Year	
	%	OR‡	(95% CI)	%	(se)	%	(se)	%	(se)
I. Anxiety disorders									
Agoraphobia	24.6§	5.5*	(4.2-7.2)	25.2	(3.7)	37.5	(7.2)	37.3	(7.6)
Panic disorder	21.9	11.3*	(8.7-14.7)	31.3	(6.1)	13.9	(6.1)	54.8	(8.6)
Simple phobia	34.1	4.7*	(3.5-6.3)	17.2	(2.6)	67.1	(5.7)	15.7	(5.5)
Social phobia	35.4	3.9*	(3.1-5.0)	27.0	(5.4)	63.2	(6.3)	9.8	(3.2)
Any anxiety disorder	60.8	5.8*	(4.8-6.9)	35.4	(3.2)	62.9	(4.6)	30.8	(4.8)
II. Mood disorders									
Dysthymia	38.9	12.6*	(9.7-16.3)	33.9	(4.4)	30.1	(4.6)	36.0	(4.3)
Major depression	61.2	9.2*	(6.8-12.6)	23.1	(4.2)	29.8	(3.0)	47.1	(3.8)
Mania	3.1	13.8*	(5.8-33.1)	28.0	(6.9)	34.2	(8.8)	37.7	(7.2)
Any mood disorder	69.9	11.8*	(9.1-15.3)	35.5	(3.5)	37.0	(2.5)	47.1	(3.3)
III. Substance use disorders									
Substance abuse	40.3	2.1*	(1.6-2.7)	47.3	(4.0)	65.5	(4.6)	12.6	(3.8)
Substance dependence	33.0	2.8*	(2.1-3.8)	51.9	(4.4)	47.9	(5.5)	16.9	(5.0)
Any substance use disorder	42.8	2.2	(1.7-2.8)	55.8	(4.0)	62.7	(4.6)	17.7	(4.1)
IV. Any disorder	90.6	11.0*	(7.7-15.8)	53.3	(3.4)	68.4	(2.0)	54.3	(3.6)

* p<0.05

† All percentages were calculated in the subsample of respondents who had both disorders.

‡ Results in this column are based on logistic regression coefficients estimated in a series of separate models to predict lifetime GAD in the United States. For example, the odds of having GAD are 5.5 times as great among respondents with a history of lifetime agoraphobia as those without lifetime agoraphobia.

§ 24.6% of people in the United States with lifetime GAD had agoraphobia in their lifetime.

|| Substances include alcohol, cocaine, inhalants, marijuana, hashish, opiates, psychedelics, sedatives, and stimulants.