

5/9/07

For table 2 (page 547), SAD should be defined in the first footnote as separation anxiety disorder, not seasonal affective disorder.

6/1/07

The tables in the paper are mislabeled and confuse the reader as to whether standard errors or standard deviations are presented. Standard deviations were requested by the editors. All tables in the paper (with the exception of table 2) present standard deviations and NOT standard errors. Table 2 was inadvertently left as standard errors and was not updated to standard deviations when we received the request from the journal. Table 2 is presented below with standard deviations rather than standard errors.

Table 2. Lifetime comorbidity of DSM-IV/CIDI bipolar disorder (BPD) with other DSM-IV/CIDI disorders in the total sample (n = 9282)

	Any BPD				BP-I				BP-II				Sub-threshold BPD				$\chi^2_{2,4}$
	% ¹	(sd)	OR ²	(95% CI)	% ¹	(sd)	OR ²	(95% CI)	% ¹	(sd)	OR ²	(95% CI)	% ¹	(sd)	OR ²	(95% CI)	
I. Anxiety disorders																	
Agoraphobia without panic	5.7	(26.4)	5.3*	(3.0-9.3)	5.6	(21.9)	5.2*	(2.1-13.0)	8.1	(28.3)	7.2*	(3.3-15.8)	4.6	(26.6)	4.3*	(1.7-10.8)	1.1
Panic disorder	20.1	(40.6)	5.8*	(4.4-7.7)	29.1	(42.0)	9.4*	(5.9-15.1)	27.2	(53.6)	8.1*	(4.7-14.1)	13.1	(36.9)	3.6*	(2.3-5.6)	13.5**
Panic attacks	61.9	(40.6)	4.3*	(3.5-5.2)	63.9	(42.9)	4.6*	(3.0-6.9)	72.9	(48.5)	7.1*	(4.3-11.5)	56.0	(45.8)	3.4*	(2.6-4.4)	6.7**
PTSD	24.2	(41.2)	4.7*	(3.3-6.8)	30.9	(33.1)	6.6*	(4.2-10.4)	34.3	(32.6)	7.3*	(4.9-10.8)	16.5	(39.0)	3.0*	(1.7-5.3)	9.1**
GAD	29.6	(50.7)	6.1*	(4.6-8.1)	38.7	(44.8)	9.4*	(6.2-14.2)	37.0	(48.5)	7.7*	(5.1-11.7)	22.3	(50.2)	4.3*	(2.8-6.7)	6.4**
Specific phobia	35.5	(56.8)	4.0*	(3.1-5.2)	47.1	(41.0)	6.5*	(4.5-9.4)	51.1	(50.6)	7.5*	(5.1-11.0)	23.3	(51.7)	2.2*	(1.5-3.4)	25.7**
Social phobia	37.8	(62.9)	4.6*	(3.5-5.9)	51.6	(56.3)	7.9*	(4.9-12.8)	54.6	(54.6)	9.0*	(6.0-13.7)	24.1	(54.7)	2.4*	(1.6-3.6)	27.8**
OCD	13.6	(29.5)	10.2*	(4.6-22.9)	25.2	(35.2)	21.4*	(7.6-60.1)	20.8	(41.8)	16.7*	(4.7-59.7)	4.3	(19.2)	3.0	(0.9-10.6)	6.6**
SAD	35.4	(31.7)	5.4*	(4.6-6.5)	41.2	(45.1)	6.8*	(4.1-11.3)	42.8	(43.7)	7.7*	(4.9-12.2)	29.4	(39.0)	4.1*	(2.9-5.7)	4.9**
Any anxiety disorder	74.9	(44.4)	6.5*	(4.7-9.0)	86.7	(29.3)	14.1*	(6.9-28.8)	89.2	(26.2)	17.6*	(9.0-34.7)	63.1	(42.5)	3.8*	(2.6-5.4)	29.9**
II. Impulse-control disorders³																	
IED	28.9	(39.6)	4.5*	(3.6-5.8)	38.1	(34.6)	6.7*	(4.4-10.1)	22.9	(38.9)	3.6*	(2.1-6.3)	27.9	(40.2)	4.2*	(2.9-6.0)	4.1
ADHD	31.4	(42.9)	6.7*	(4.8-9.3)	40.6	(38.9)	10.0*	(6.1-16.1)	42.3	(46.8)	10.4*	(5.7-18.8)	23.0	(38.5)	4.4*	(2.6-7.4)	7.4**
ODD	36.8	(41.6)	6.1*	(4.5-8.4)	44.4	(42.2)	8.5*	(5.1-14.1)	38.2	(45.0)	7.0*	(3.5-14.3)	32.8	(40.5)	5.0*	(3.5-7.1)	3.3
CD	30.3	(41.6)	4.9*	(3.6-6.7)	43.8	(38.9)	8.9*	(5.6-14.2)	18.6	(31.9)	2.6*	(1.3-5.1)	28.9	(37.5)	4.5*	(3.0-6.8)	10.6**
Any impulse-control disorder	62.8	(38.9)	5.6*	(4.3-7.3)	71.2	(33.6)	8.3*	(5.2-13.2)	70.4	(41.8)	8.1*	(4.3-15.4)	56.1	(37.5)	4.2*	(3.0-6.0)	7.5**
III. Substance use disorders																	
Alcohol abuse	39.1	(41.2)	4.3*	(3.3-5.5)	56.3	(27.1)	8.6*	(5.9-12.4)	36.0	(34.9)	3.7*	(2.4-5.7)	33.2	(49.4)	3.3*	(2.3-4.8)	17.2**
Alcohol dependence	23.2	(30.1)	5.7*	(4.3-7.6)	38.0	(37.6)	11.6*	(6.8-19.7)	19.0	(30.2)	4.4*	(2.4-7.8)	18.9	(40.2)	4.5*	(3.0-6.8)	10.2**
Drug abuse	28.8	(42.8)	4.5*	(3.3-5.9)	48.3	(28.6)	10.1*	(7.0-14.4)	23.7	(27.0)	3.5*	(2.4-5.3)	22.9	(44.8)	3.2*	(2.1-5.1)	15.4**
Drug dependence	14.0	(28.5)	5.2*	(3.7-7.2)	30.4	(30.8)	13.7*	(8.9-21.1)	8.7	(24.6)	3.1*	(1.4-6.8)	9.5	(26.4)	3.4*	(2.0-5.7)	18.3**
Any substance	42.3	(42.8)	4.2*	(3.3-5.5)	60.3	(31.6)	8.8*	(5.9-13.1)	40.4	(29.4)	3.9*	(2.7-5.7)	35.5	(49.4)	3.2*	(2.2-4.6)	18.8**
IV. Any disorder																	
Any disorder ⁴	92.3	(29.5)	13.1*	(6.7-25.5)	97.7	(10.6)	47.7*	(11.3-201.3)	95.8	(15.6)	24.5*	(7.1-85.1)	88.4	(36.6)	8.5*	(3.9-18.2)	10.7**
Exactly 1 disorder	12.7	(26.8)	4.8*	(2.2-10.4)	8.1	(20.4)	10.2*	(1.9-54.3)	7.0	(16.2)	4.8*	(1.2-19.0)	17.1	(36.6)	4.3*	(1.7-10.9)	0.8
Exactly 2 disorders	9.4	(22.8)	5.6*	(2.5-12.5)	3.4	(13.8)	6.9*	(1.0-48.5)	2.9	(13.7)	3.3	(0.5-22.8)	14.7	(26.7)	5.8*	(2.3-14.7)	0.3
3 or more disorders	70.1	(33.5)	26.4*	(13.7-50.8)	86.2	(24.4)	112.3*	(27.2-464.6)	85.8	(26.2)	58.3*	(16.3-208.4)	56.7	(37.5)	14.3*	(6.9-29.5)	18.1**

*Significant at the .05 level, two-sided test

**Significant difference across the three BPD subgroups at the .05 level, two-sided test

¹The prevalence of the comorbid disorder among respondents with BPD

²Based on logistic regression models with one DSM-IV/CIDI disorder at a time as a predictor of LT BPD, controlling for age at interview (5-year intervals) sex, and race-ethnicity. The last model had predictors

“exactly one,” “exactly two” and “exactly three” disorders in one model.

³Sample included Part II respondents whose age at interview was less than or equal to 44 (n = 3197)

⁴Test of the significance of difference in ORs across the three BPD subgroups