

**Appendix Table 1. Numbers of respondents by age group in each country<sup>1</sup>**

	All ages			18-34			Age Groups 35-49			50-64			65+			Age Difference
	n <sup>4</sup>	%	(sd)	n <sup>4</sup>	%	(sd)	n <sup>4</sup>	%	(sd)	n <sup>4</sup>	%	(sd)	n <sup>4</sup>	%	(sd)	$\chi^2_3$
<b>I. Developed</b>																
Belgium	2419	5.0	5.3	573	6.1	10.0	775	5.2	5.9	570	5.8	7.3	501	2.4	2.4	9.1*
France	2894	5.9	7.9	743	8.3	10.4	942	6.0	7.4	719	5.6	7.3	490	2.5	2.9	19.1*
Germany	3555	3.0	3.1	815	4.7	5.1	1180	2.8	2.9	893	2.9	3.6	667	1.3	1.2	13.5*
Israel <sup>2</sup>	4859	6.1	6.8	1627	6.3	6.1	1302	5.7	6.0	1069	6.2	6.3	861	6.0	6.3	0.5
Italy	4712	3.0	2.3	1326	2.8	2.4	1393	2.3	2.1	1153	3.8	3.9	840	3.3	3.4	6.2
Japan	4129	2.2	3.1	683	4.0	4.5	976	2.1	2.2	1268	2.1	2.2	1202	1.0	1.1	10.8*
Netherlands	2372	4.9	5.8	564	5.9	7.2	729	5.7	8.2	627	4.3	5.1	452	2.4	2.5	9.2*
New Zealand <sup>2</sup>	12992	6.6	9.0	1333	9.9	13.2	5304	7.6	10.7	3909	5.8	7.9	2244	1.8	2.2	138.1*
Spain	5473	4.0	4.7	1567	3.6	5.5	1431	3.8	3.9	1024	5.4	5.7	1451	3.6	4.6	6.5
United States	9282	8.3	8.5	3033	10.4	9.0	2865	9.4	8.4	1922	7.7	8.9	1461	2.6	2.6	103.5*
Combined	51771	5.5	5.5	14561	7.0	9.9	15532	6.0	6.3	11736	5.1	5.0	9942	2.6	3.3	150.2*
<b>II. Developing</b>																
Brazil	5037	10.4	13.9	1841	10.9	9.9	1729	11.8	17.7	1045	9.1	11.2	422	3.9	5.0	36.1*
China-Shenzhen	7134	3.8	4.7	4565	4.3	4	2010	2.4	3	403	1.3	1.4	154	2.3	1.4	26.1*
Colombia <sup>3</sup>	4426	6.2	6.7	1431	7.6	9.0	1735	5.5	6.9	730	4.8	6.4	530	5.9	6.3	5.3
India	2992	4.5	4.9	1174	2.2	1.7	1035	6.3	6.6	542	7.9	7.6	241	5.2	5.5	33.0*
Lebanon	2857	5.5	8.4	965	5.2	7.5	931	6.6	8.2	553	5.8	5.6	408	3.1	3.6	6.1
Mexico <sup>3</sup>	5782	4.0	4.9	2060	3.7	4.6	2236	3.7	4.0	840	4.6	5.9	646	5.2	5.2	6.3
South Africa	4315	4.9	6.0	2172	4.5	6.0	1264	5.4	7.7	638	5.6	6.2	241	3.5	3.8	3
Ukraine	4724	8.4	12.5	1193	6.0	6.1	1225	6.7	7.5	1180	10.1	12.3	1126	13.0	20.3	21.2*
Combined	37265	5.9	9.6	16781	5.3	6.0	11659	6.2	8.1	6112	6.8	8.5	2713	7.5	11.5	2.2

These are the denominators (unweighted) on which the prevalence estimates in Table 2 of the paper were calculated.

<sup>1</sup>The numbers reported in this table are the unweighted number of respondents in the samples in each survey. These are the denominators on which the prevalence estimates in Table 2 in the paper were based.

<sup>2</sup>The lower end of the age range was 16 in New Zealand and 21 in Israel.

<sup>3</sup>The upper end of the age range was 65 in Colombia and Mexico. As a result, the age groups used in these countries was 18-29, 30-44, 45-54, 55+

<sup>4</sup>Denominator n for the %s reported. Note that some countries had different cutpoints for age groups (see first and second footnote), thus the Ns at country level may not add up to the combined denominator n.