Appendix Table 1. Multivariate socio-demographic predictors of 12-month treatment and adequacy among respondents with 12-month CIDI/DSM-IV MDD, controlling for clinical

predictors (n=514)¹

predicto	ors (II-514)	Any OR	Treatment (95% CI)	Adequ	(95% CI)
	18-29 30-44 45-59 60+	2.1 2.2 4.0 1.0	(0.5-8.5) (0.6-7.8) (1.2-13.6) χ^2_3 =8.9*	2.7 2.2 2.5 1.0	(0.9-8.3) (0.5-9.6) (0.5-12.5) χ^2_3 =4.1
	Female Male	1.4 1.0	$\begin{array}{c} (0.7-2.5) \\ \\ \chi^2_1 = 1.0 \end{array}$	0.6 1.0	$\begin{array}{c} (0.3-1.1) \\ - \\ \chi^2_1 = 2.9 \end{array}$
	hnicity Hispanic Non-Hispanic Black Non-Hispanic White Other	0.4 0.6 1.0 0.2	$\begin{array}{c} (0.2\text{-}1.3) \\ (0.3\text{-}1.3) \\ - \\ (0.1\text{-}0.8) \\ \chi^2_3 = 10.9^* \end{array}$	0.9 1.1 1.0 0.2	$\begin{array}{c} (0.3-2.5) \\ (0.5-2.5) \\ \\ (0.1-1.2) \\ \chi^2_3 = 3.5 \end{array}$
	ment status Employed Homemaker Retired Student Other	1.0 1.1 1.0 1.4 1.7	$\begin{array}{c} -0.4 - 3.0 \\ (0.4 - 3.0) \\ (0.3 - 3.6) \\ (0.5 - 4.3) \\ (0.8 - 3.9) \\ \chi^2_4 = 1.9 \end{array}$	1.0 0.5 0.7 1.9 0.6	$\begin{array}{c} -2 \\ (0.2 - 1.4) \\ (0.2 - 2.5) \\ (0.6 - 5.8) \\ (0.3 - 1.4) \\ \chi^{2}_{4} = 4.3 \end{array}$
	Married Never Married Divorced/Separated/Widowed	1.0 1.3 1.4	$\begin{array}{c} -1.3 \\ -2.3 \\ (0.7-2.3) \\ (0.9-2.3) \\ \chi^2_2 = 2.2 \end{array}$	1.0 0.8 0.9	$\begin{array}{c} -2.3 \\ -2.3 \\ (0.3-1.9) \\ (0.5-1.7) \\ \chi^{2}_{2}=0.3 \end{array}$
	on 0-11 12 13-15 16+	0.4 0.7 0.4 1.0	$\begin{array}{c} (0.2\text{-}0.8) \\ (0.4\text{-}1.2) \\ (0.2\text{-}0.7) \\ \\ \chi^2_3 = 17.0^* \end{array}$	0.7 0.4 0.8 1.0	$(0.3-1.7)$ $(0.2-0.8)$ $(0.5-1.4)$ $-$ $\chi^{2}_{3}=7.3$
	Categories Below Poverty 1-3 Times Poverty 3-6 Times Poverty 6+ Times Poverty	0.6 0.6 0.9 1.0	$\begin{array}{c} (0.3-1.3) \\ (0.3-1.2) \\ (0.4-1.8) \\ - \\ \chi^2_3 = 3.2 \end{array}$	0.9 0.6 1.1 1.0	$(0.3-2.2)$ $(0.2-1.3)$ $(0.6-1.9)$ $-$ $\chi^{2}_{3}=3.0$
	Northeast Midwest South West	1.3 1.0 1.0 0.9	$\begin{array}{c} (0.6\text{-}3.1) \\ (0.5\text{-}2.0) \\ - \\ (0.5\text{-}1.7) \\ \chi^2_3 = 0.8 \end{array}$	1.4 1.4 1.0 1.3	$(0.7-2.9)$ $(0.7-2.6)$ $(0.7-2.4)$ $\chi^{2}_{3}=1.5$
	ty Major Metropolitan Other Urban Rural	1.8 1.4 1.0	$\begin{array}{c} (0.8\text{-}4.0) \\ (0.7\text{-}2.8) \\ \\ \chi^2_2 = 2.4 \end{array}$	1.0 0.8 1.0	$\begin{array}{c} (0.5\text{-}2.0) \\ (0.4\text{-}1.6) \\ \\ \chi^2_2 = 0.7 \end{array}$

^{*} Significant at the .05 level, two-sided test.

¹ Clinical predictors include all the predictors in Table 9. Cases exclude respondents who had missing or incomplete QIDS scores or who scored in the QIDS non-case range.