

Appendix Table 1. AUC and stratum-specific PPV of the CIDI-based BPD screening scale for DSM-IV BP-I/II based on endorsement of the CIDI euphoria diagnostic stem question in the total sample and a number of critical sub-samples

Imputation stratum	The sample in which PPV is calculated											
	Total		High PCP ¹		WT PCP ¹		SMH ¹		EMPL ¹		LOW INC ¹	
	PPV	(se)	PPV	(se)	PPV	(se)	PPV	(se)	PPV	(se)	PPV	(se)
Negative Criterion B screen ²	0.1	(0.0)	0.2	(0.2)	0.3	(0.1)	0.7	(0.3)	0.1	(0.1)	0.2	(0.1)
Number Endorsed/Positive Criterion B screen ²												
0	0.3	(0.3)	0.0	(0.0)	0.3	(0.3)	0.0	(0.0)	0.4	(0.4)	0.0	(0.0)
1	0.3	(0.3)	0.0	(0.0)	0.3	(0.3)	0.0	(0.0)	0.4	(0.4)	0.0	(0.0)
2	0.3	(0.3)	0.0	(0.0)	0.3	(0.3)	0.0	(0.0)	0.4	(0.4)	0.0	(0.0)
3	3.3	(1.1)	4.9	(4.8)	3.5	(1.7)	13.8	(6.0)	2.4	(0.9)	0.0	(0.0)
4	3.3	(1.1)	4.9	(4.8)	3.5	(1.7)	13.8	(6.0)	2.4	(0.9)	0.0	(0.0)
5	22.0	(3.5)	18.4	(8.6)	22.0	(4.7)	42.0	(8.9)	18.8	(4.3)	17.4	(7.4)
6	31.0	(4.3)	42.5	(12.9)	42.9	(11.9)	48.2	(9.2)	26.6	(4.8)	40.5	(5.9)
7	41.5	(5.8)	52.0	(9.4)	52.6	(8.3)	55.2	(9.5)	39.5	(5.3)	40.5	(5.9)
8	56.3	(6.7)	52.0	(9.4)	52.6	(8.3)	69.6	(11.1)	52.0	(9.0)	40.5	(5.9)
9	67.5	(14.9)	100.0	(0.0)	85.3	(8.8)	88.2	(11.2)	57.7	(18.2)	50.3	(22.2)
AUC	.844		.845		.841		.774		.849		.821	
(n)	(9282)		(578)		(5225)		(821)		(6314)		(695)	

¹High PCP = respondents who saw a primary care physician at least 12 times in the year before interview; WT PCP = respondents who was a primary care physician at least ones in the year before interview weighted for number of visits; SMH = respondents who received specialty mental health treatment in the year before interview; LOW INC = respondents with family incomes at or below the federal poverty line.

²Negative stem = respondents who either failed to endorse the CIDI BPD euphoria diagnostic stem question or endorsed the diagnostic stem question but failed to endorse the Criterion B screening question; Number endorsed/Positive Criterion B screen = number of Criterion B symptom questions endorsed out of a maximum of nine among respondents who endorsed the Criterion B screening question. Equivalence of PPV for some numbers is due to these sub-samples being collapsed because of insignificant differences in PPV.

Appendix Table 2. AUC and stratum-specific PPV of the CIDI-based BPD screening scale for DSM-IV BP-/II based on endorsement of the CIDI euphoria or irritability diagnostic stem questions in the total sample and a number of critical sub-samples

Imputation stratum	The sample in which PPV is calculated											
	Total		High PCP ¹		WT PCP ¹		SMH ¹		EMPL ¹		LOW INC ¹	
	PPV	(se)	PPV	(se)	PPV	(se)	PPV	(se)	PPV	(se)	PPV	(se)
Negative Criterion B screen ²	0.0	(0.0)	0.0	(0.0)	0.0	(0.0)	0.0	(0.0)	0.0	(0.0)	0.0	(0.0)
Number Endorsed/Positive Criterion B screen ²												
0	0.2	(0.2)	0.0	(0.0)	0.2	(0.2)	0.0	(0.0)	0.3	(0.3)	0.0	(0.0)
1	0.2	(0.2)	0.0	(0.0)	0.2	(0.2)	0.0	(0.0)	0.3	(0.3)	0.0	(0.0)
2	0.2	(0.2)	0.0	(0.0)	0.2	(0.2)	0.0	(0.0)	0.3	(0.3)	0.0	(0.0)
3	3.0	(0.9)	3.6	(3.5)	3.0	(1.3)	10.4	(4.7)	2.2	(0.8)	0.0	(0.0)
4	3.0	(0.9)	3.6	(3.5)	3.0	(1.3)	10.4	(4.7)	2.2	(0.8)	0.0	(0.0)
5	19.7	(3.2)	17.0	(8.1)	20.8	(4.3)	39.0	(6.3)	16.0	(3.8)	15.5	(6.7)
6	26.8	(3.5)	33.4	(11.0)	37.2	(10.8)	39.0	(6.3)	24.5	(4.2)	31.1	(8.7)
7	39.5	(5.9)	52.6	(14.9)	50.2	(10.9)	55.2	(9.5)	37.2	(5.4)	36.1	(10.9)
8	56.2	(6.2)	54.9	(14.2)	53.7	(12.0)	71.0	(10.9)	51.7	(8.3)	44.2	(10.1)
9	67.5	(14.9)	100.0	(0.0)	85.3	(8.8)	88.2	(11.2)	57.7	(18.2)	50.3	(22.2)
AUC	.852		.865		.854		.800		.858		.848	
(n)	(9282)		(578)		(5225)		(821)		(6314)		(695)	

¹High PCP = respondents who saw a primary care physician at least 12 times in the year before interview; WT PCP = respondents who was a primary care physician at least ones in the year before interview weighted for number of visits; SMH = respondents who received specialty mental health treatment in the year before interview; LOW INC = respondents with family incomes at or below the federal poverty line.

²Negative stem = respondents who either failed to endorse the CIDI BPD euphoria diagnostic stem question or endorsed the diagnostic stem question but failed to endorse the Criterion B screening question; Number endorsed/Positive Criterion B screen = number of Criterion B symptom questions endorsed out of a maximum of nine among respondents who endorsed the Criterion B screening question. Equivalence of PPV for some numbers is due to these sub-samples being collapsed because of insignificant differences in PPV.

Appendix Table 3. AUC and stratum-specific PPV of the CIDI-based BPD screening scale for DSM-IV BP-I/II/sub based on endorsement of the CIDI euphoria diagnostic stem question in the total sample and a number of critical sub-samples

Imputation stratum	The sample in which PPV is calculated											
	Total		High PCP ¹		WT PCP ¹		SMH ¹		EMPL ¹		LOW INC ¹	
	PPV	(se)	PPV	(se)	PPV	(se)	PPV	(se)	PPV	(se)	PPV	(se)
Negative Criterion B screen ²	0.7	(0.1)	2.0	(0.9)	1.2	(0.3)	2.8	(0.8)	0.8	(0.2)	0.7	(0.2)
Number Endorsed/Positive Criterion B screen ²												
0	3.2	(1.6)	0.0	(0.0)	2.0	(1.4)	0.0	(0.0)	4.5	(2.2)	10.5	(7.7)
1	3.2	(1.6)	0.0	(0.0)	2.0	(1.4)	0.0	(0.0)	4.5	(2.2)	10.5	(7.7)
2	9.3	(3.1)	15.5	(9.8)	11.6	(5.1)	0.0	(0.0)	8.5	(3.4)	18.2	(6.0)
3	23.6	(4.2)	25.1	(13.3)	26.4	(5.3)	33.1	(11.0)	25.3	(5.2)	18.2	(6.0)
4	32.1	(3.9)	25.8	(11.3)	40.3	(11.7)	53.5	(9.9)	33.7	(4.8)	18.2	(6.0)
5	44.2	(3.4)	43.3	(7.4)	47.7	(6.2)	60.1	(5.2)	40.8	(3.1)	47.4	(6.6)
6	44.2	(3.4)	43.3	(7.4)	47.7	(6.2)	60.1	(5.2)	40.8	(3.1)	47.4	(6.6)
7	57.0	(6.1)	55.2	(14.7)	61.2	(10.4)	60.1	(5.2)	56.2	(6.0)	47.4	(6.6)
8	67.2	(6.9)	70.4	(12.5)	66.1	(12.5)	76.5	(9.4)	63.2	(9.2)	57.7	(12.1)
9	67.2	(6.9)	70.4	(12.5)	66.1	(12.5)	76.5	(9.4)	63.2	(9.2)	57.7	(12.1)
AUC	.744		.736		.732		.699		.730		.718	
(n)	(9282)		(578)		(5225)		(821)		(6314)		(695)	

¹High PCP = respondents who saw a primary care physician at least 12 times in the year before interview; WT PCP = respondents who was a primary care physician at least ones in the year before interview weighted for number of visits; SMH = respondents who received specialty mental health treatment in the year before interview; LOW INC = respondents with family incomes at or below the federal poverty line.

²Negative stem = respondents who either failed to endorse either of the two CIDI BPD diagnostic stem questions or endorsed a diagnostic stem question but failed to endorse the Criterion B screening question; Number endorsed/Positive Criterion B screen = number of Criterion B symptom questions endorsed out of a maximum of nine among respondents who endorsed the Criterion B screening question. Equivalence of PPV for some numbers is due to these sub-samples being collapsed because of insignificant differences in PPV.

Appendix Table 4. AUC and stratum-specific PPV of the CIDI-based BPD screening scale for DSM-IV BP-I/II/sub based on endorsement of either the CIDI euphoria or irritability diagnostic stem questions in the total sample and a number of critical sub-samples

Imputation stratum	The sample in which PPV is calculated											
	Total		High PCP ¹		WT PCP ¹		SMH ¹		EMPL ¹		LOW INC ¹	
	PPV	(se)	PPV	(se)	PPV	(se)	PPV	(se)	PPV	(se)	PPV	(se)
Negative Criterion B screen ²	0.0	(0.0)	0.0	(0.0)	0.0	(0.0)	0.0	(0.0)	0.0	(0.0)	0.0	(0.0)
Number Endorsed/Positive Criterion B screen ²												
0	2.5	(1.3)	0.0	(0.0)	1.7	(1.1)	0.0	(0.0)	3.5	(1.8)	7.5	(5.4)
1	2.5	(1.3)	0.0	(0.0)	1.7	(1.1)	0.0	(0.0)	3.5	(1.8)	7.5	(5.4)
2	7.2	(2.3)	9.1	(6.3)	7.1	(3.7)	0.0	(0.0)	7.1	(2.7)	10.5	(5.1)
3	21.5	(3.7)	29.6	(8.5)	25.7	(5.0)	30.4	(9.9)	23.4	(4.5)	10.5	(5.1)
4	30.6	(3.1)	29.6	(8.5)	37.1	(10.4)	45.4	(8.6)	31.7	(3.9)	21.2	(7.2)
5	46.2	(3.1)	44.0	(6.2)	47.2	(5.6)	62.3	(4.6)	43.8	(3.2)	45.6	(6.1)
6	46.2	(3.1)	44.0	(6.2)	47.2	(5.6)	62.3	(4.6)	43.8	(3.2)	45.6	(6.1)
7	58.4	(5.7)	55.2	(14.7)	61.4	(9.6)	62.3	(4.6)	56.6	(5.8)	45.6	(6.1)
8	68.6	(6.5)	72.1	(11.9)	67.4	(11.8)	77.4	(9.0)	65.3	(8.3)	59.8	(11.7)
9	68.6	(6.5)	72.1	(11.9)	67.4	(11.8)	77.4	(9.0)	65.3	(8.3)	59.8	(11.7)
AUC	.765		.746		.755		.738		.750		.739	
(n)	(9282)		(578)		(5225)		(821)		(6314)		(695)	

¹High PCP = respondents who saw a primary care physician at least 12 times in the year before interview; WT PCP = respondents who was a primary care physician at least ones in the year before interview weighted for number of visits; SMH = respondents who received specialty mental health treatment in the year before interview; LOW INC = respondents with family incomes at or below the federal poverty line.

²Negative stem = respondents who either failed to endorse either of the two CIDI BPD diagnostic stem questions or endorsed a diagnostic stem question but failed to endorse the Criterion B screening question; Number endorsed/Positive Criterion B screen = number of Criterion B symptom questions endorsed out of a maximum of nine among respondents who endorsed the Criterion B screening question. Equivalence of PPV for some numbers is due to these sub-samples being collapsed because of insignificant differences in PPV.