

4/6/07

The prevalence rate of anorexia is 0.5 percent, not 0.6 percent as reported on page 350 of this publication (in both the results section and in table 1).

4/20/12

It has come to our attention that the last row of table 7 is incorrect in the published paper. We have revised the estimates for the prevalence of treatment specifically for eating disorders among 12-month cases, and the updated table is below. In the published table this row mistakenly reported the prevalence of any treatment for emotional problems among 12-month cases.

Table 7. Lifetime and 12-month treatment of DSM-IV eating disorders

	Anorexia Nervosa		Bulimia Nervosa		Binge-eating Disorder	
	%	(se)	%	(se)	%	(se)
I. Lifetime treatment¹ for any emotional problem						
General medical	45.3	(12.6)	43.9	(11.4)	36.3	(5.6)
Psychiatrist	29.4	(12.8)	48.2	(8.1)	33.6	(6.3)
Other mental health	34.3	(10.2)	48.3	(9.9)	35.0	(3.2)
Human services	11.7	(7.5)	14.9	(3.0)	19.9	(4.2)
CAM	26.0	(12.1)	21.6	(6.1)	19.7	(4.1)
Any lifetime treatment	50.0	(14.6)	63.2	(7.8)	51.2	(6.5)
(n)	(23)		(52)		(115)	
II. Twelve-month treatment¹ for any emotional problem²						
General medical			26.1	(14.3)	20.8	(7.2)
Psychiatrist			9.0	(5.5)	5.0	(2.6)
Other mental health			3.7	(3.3)	24.0	(6.3)
Human services			13.5	(10.0)	7.1	(4.6)
CAM			0.0	--	12.4	(5.1)
Any 12-month treatment			15.6	(9.4)	28.5	(10.7)
(n)			(16)		(51)	
III. Treatment of eating disorders						
Lifetime	33.8	(14.2)	43.2	(9.7)	43.6	(6.2)
Twelve-month			10.3	(5.1)	6.4	(2.9)

¹General medical treatment is treatment by a non-psychiatrist physician or nurse or other medical practitioner who is not a mental health specialist. Psychiatrist treatment is treatment by a psychiatrist. Other mental health treatment is treatment by any mental health specialist other than a psychiatrist (e.g., psychologist, psychiatric social worker). Human services treatment is treatment by a minister, priest, rabbi, or other spiritual advisor or by a caseworker in a social services agency. CAM (Complementary-alternative medical) treatment is treatment in a self-help group or treatment by an alternative medical provider (e.g., massage therapist, chiropractor).

²There were no respondents with 12-month Anorexia Nervosa.