

Appendix Table 2. Associations between Lifetime Anxiety Disorders and Lifetime Major Depressive Episodes in the ICPE Surveys

| Brazil | Canada | Chile | Czech Republic | Germany | Japan | Mexico | Netherlands | Turkey | U.S. | |
|---|------------------|-------------------|-----------------|-------------------|----------------|-----------------|-----------------|-----------------|-------------------|-----------------|
| I. Bivariate associations (odds ratios) between lifetime anxiety disorders and lifetime major depressive episodes | | | | | | | | | | |
| OR (95% CI) | OR (95% CI) | OR (95% CI) | OR (95% CI) | OR (95% CI) | OR (95% CI) | OR (95% CI) | OR (95% CI) | OR (95% CI) | OR (95% CI) | |
| AGOR ^a | 3.2* (1.8-5.4) | 5.2* (3.4-7.9) | 2.2* (1.4-3.6) | 7.3* (2.3-23.4) | 2.5* (1.4-4.4) | --- | 6.2* (3.6-10.6) | 4.3* (3.2-5.7) | 8.1* (5.3-12.2) | 4.4* (3.3-5.8) |
| GAD ^a | 17.2* (7.3-40.4) | 20.7* (14.4-29.7) | 6.3* (3.6-11.1) | --- | 3.0* (1.1-8.3) | 5.7* (1.6-20.7) | 3.5 (0.7-18.0) | 9.7* (7.8-12.1) | --- | 9.2* (6.8-12.6) |
| OCD ^a | 8.5* (2.0-36.2) | --- | 3.9* (3.5-4.4) | 15.8* (2.0-122.2) | 1.5 (0.5-4.5) | --- | --- | 8.0* (4.4-14.3) | 18.8* (10.6-33.3) | --- |
| PD ^a | 9.2* (3.6-23.6) | 10.5* (7.4-14.7) | 4.5* (2.4-8.5) | --- | 4.3* (2.1-9.1) | 7.3 (0.7-73.8) | --- | 8.7* (6.3-12.1) | 23.9* (10.5-54.1) | 6.8* (5.1-9.0) |
| PTSD ^a | --- | --- | 2.3* (1.1-4.6) | 8.5* (3.1-23.4) | 4.8* (2.3-9.9) | --- | --- | --- | --- | 5.7* (4.5-7.2) |
| SiP ^a | 3.5* (2.1-6.0) | 6.3* (4.5-8.7) | 4.9* (3.9-6.1) | 4.2* (2.5-7.0) | 2.8* (1.4-5.3) | --- | 8.3* (3.8-18.4) | 4.4* (3.7-5.1) | 7.8* (5.6-11.0) | 4.2* (3.5-5.1) |
| SoP ^a | 3.0* (1.8-5.1) | 3.3* (2.7-4.0) | 5.7* (4.4-7.3) | --- | --- | --- | 5.5* (2.5-12.0) | 5.3* (4.3-6.6) | 9.6* (6.8-13.6) | 3.6* (3.0-4.4) |
| Any ^a | 4.9* (3.2-7.5) | 4.0* (3.0-5.3) | 3.2* (2.0-4.9) | 5.2* (3.1-8.6) | 3.9* (2.7-5.5) | 6.3* (2.0-20.0) | 6.9* (4.4-10.8) | 6.3* (5.3-7.4) | 10.3* (8.1-13.1) | 5.3* (4.3-6.5) |
| II. Lifetime prevalences of anxiety disorders among respondents with lifetime major depressive episodes | | | | | | | | | | |
| % (se) | % (se) | % (se) | % (se) | % (se) | % (se) | % (se) | % (se) | % (se) | % (se) | |
| AGOR ^a | 12.6 (2.7) | 20.6 (2.9) | 20.9 (3.5) | 5.5 (2.3) | 6.4 (1.2) | --- | 16.3 (3.2) | 9.1 (1.0) | 9.6 (1.2) | 17.0 (1.4) |
| GAD ^a | 13.7 (3.1) | 14.3 (1.8) | 10.3 (2.0) | --- | 1.9 (0.8) | 0.1 (0.0) | 2.7 (1.8) | 20.1 (1.2) | 100.0 (0.0) | 18.2 (1.4) |
| OCD ^a | 2.8 (1.4) | --- | 3.4 (1.8) | 1.9 (1.1) | 1.0 (0.5) | --- | --- | 3.3 (0.6) | 6.5 (0.9) | --- |
| PD ^a | 5.8 (1.6) | 9.2 (1.2) | 5.3 (1.8) | 0.0 (0.0) | 4.9 (1.3) | 0.0 (0.0) | --- | 13.4 (1.4) | 4.4 (0.9) | 11.2 (1.0) |
| PTSD ^a | --- | --- | 8.7 (2.2) | 5.6 (1.9) | 4.3 (1.2) | --- | --- | --- | --- | 21.1 (1.1) |
| SiP ^a | 20.1 (3.6) | 31.8 (3.0) | 28.5 (3.7) | 20.0 (3.1) | 5.2 (1.1) | --- | 14.0 (3.5) | 25.3 (1.2) | 17.1 (1.6) | 26.7 (1.8) |
| SoP ^a | 14.7 (2.7) | 30.8 (2.1) | 31.9 (7.3) | --- | --- | --- | 9.0 (3.1) | 22.3 (1.3) | 16.3 (1.4) | 29.3 (2.0) |
| Any ^a | 44.3 (3.9) | 48.3 (3.8) | 39.1 (6.0) | 27.6 (3.7) | 19.2 (2.0) | 11.4 (0.4) | 31.1 (4.1) | 51.5 (1.5) | 38.6 (1.8) | 60.1 (2.3) |
| (n) | (1,464) | (6,902) | (2,978) | (1,534) | (3,021) | (1,029) | (1,734) | (7,076) | (6,095) | (5,877) |

^a AGOR = agoraphobia with or without panic disorder; GAD = generalized anxiety disorder; OCD = obsessive-compulsive disorder; PD = panic disorder; PTSD = posttraumatic stress disorder

SiP = simple phobia; SoP = social phobia; and "Any" = any of the anxiety disorders assessed in the survey.

* Significant at the .05 level, two-sided test