Appendix Table 8. The Effects of Temporally Primary Posttraumatic Stress Disorder (PTSD) in Predicting

| | Current | | | | Past | | | | |
|----------------|---------|-------------|-----------------|-----------|------|-----------|-------------|-----------|----------------|
| | Current | | Number of Years | | Past | | Years Since | | χ ² |
| | OR | (95% CI) | OR | (95% CI) | OR | (95% CI) | OR | (95% CI) | |
| Brazil | | | | | | | | | |
| Canada | | | | | | | | | |
| Chile | 3.6 | (0.9-14.5) | 0.9 | (0.8-1.1) | 0.2 | (0.0-1.7) | 1.2 | (1.0-1.4) | 14.5** |
| Czech Republic | 23.6* | (5.5-101.4) | | | | | 1.4* | (1.2-1.5) | 44.0** |
| Germany | 4.3* | (1.1-17.3) | 1.0 | (0.8-1.2) | | | 1.1 | (0.6-1.7) | 8.4** |
| Japan | | | | | | | | | |
| Mexico | | | | | | | | | |
| Netherlands | | | | | | | | | |
| Turkey | | | | | | | | | |
| U.S. | 10.3* | (6.4-16.5) | 0.9* | (0.8-0.9) | 1.1 | (0.4-3.1) | 1.1 | (1.0-1.3) | 193.7** |

First Onset of Major Depressive Episodes, Separately by Country

* Significant at the .05 level, two-sided test

** Significant at the .05 level