Appendix Table 1. Multivariate associations (odds ratios) between childhood SES and the persistence of DSM-IV/CIDI disorders¹

	Any Disorder		Mood		Anxiety		Substance		Behavior	
	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)
I. Without CA controls ² Parent Occupation										
Never worked/Elementary	1.1	(0.9- 1.3)	1.0	(0.8- 1.3)	1.1	(0.9- 1.5)	1.1	(0.8- 1.6)	0.8	(0.5- 1.2)
Low-average prestige	1.0	(0.9- 1.2)	1.0	(0.8- 1.2)	1.1	(0.9- 1.3)	1.2	(1.0- 1.4)	0.9	(0.7- 1.1)
High-average prestige	1.0	(0.9- 1.1)	8.0	(0.6- 1.0)	0.9	(0.8- 1.1)	1.3	(1.0- 1.7)	1.0	(0.8- 1.4)
High prestige	-		-		-		-		-	
χ^2 3		1.0		4.0		3.8		4.1		3.6
Parent Education										
0-11 years	1.1	(0.9- 1.3)	1.2	(0.9- 1.6)	1.0	(0.8- 1.3)	1.0	(0.8- 1.2)	1.6*	(1.0- 2.5)
12 years	1.1	(1.0- 1.2)	1.3	(1.0- 1.6)	1.2	(1.0- 1.3)	1.0	(0.8- 1.2)	0.9	(0.7- 1.3)
13-15 years	1.1	(0.9- 1.2)	1.2	(0.9- 1.5)	1.0	(0.8- 1.2)	1.0	(0.8- 1.4)	1.2	(0.9- 1.6)
16+ years	-		-		-		-		-	
χ^2_3		1.6		5.1		4.9		0.3		9.6*
Financial Hardship										
Yes	1.1	(0.9- 1.2)	1.2	(0.9- 1.5)	1.1	(0.9- 1.3)	1.0	(0.8- 1.2)	1.2	(0.9- 1.6)
χ^2_1		0.9		1.4		0.9		0.2		1.1
II. With CA controls ³										
Parent Occupation										
Never worked/Elementary	1.1	(0.9- 1.3)	1.0	(0.8- 1.4)	1.1	(0.8- 1.4)	1.2	(0.8- 1.7)	8.0	(0.5- 1.2)
Low-average prestige	1.0	(0.9- 1.2)	1.0	(0.9- 1.3)	1.0	(0.9- 1.3)	1.2	(1.0- 1.4)	8.0	(0.6- 1.0)
High-average prestige	1.0	(0.9- 1.1)	0.8	(0.6- 1.0)	0.9	(0.8- 1.1)	1.2	(0.9- 1.6)	1.0	(0.8- 1.4)
High prestige	-		-		-		-		-	
χ^2 3		1.0		4.8		3.9		4.6		5.6
Parent Education										
0-11 years	1.1	(0.9- 1.3)	1.2	(0.9- 1.6)	1.0	(0.8- 1.3)	1.0	(0.8- 1.2)	1.6*	(1.0- 2.7)
12 years	1.1	(1.0- 1.2)	1.3	(1.0- 1.6)	1.2	(1.0- 1.3)	0.9	(0.7- 1.1)	1.0	(0.8- 1.3)
13-15 years	1.1	(1.0- 1.2)	1.2	(0.9- 1.5)	1.0	(0.8- 1.2)	1.0	(0.8- 1.4)	1.4*	(1.1- 1.8)
16+ years	-		-		-		-		-	
χ^2_3		1.9		4.4		5.3		1.0		12.1*
Financial Hardship										
Yes	1.0	(0.9- 1.1)	1.1	(0.9- 1.4)	1.0	(0.8- 1.2)	0.9	(0.7- 1.2)	0.9	(0.7- 1.2)
χ^2 1 .		0.1		0.9		0.0		0.4	0.4	
$(n_1/n_2)^4$	(7,6	07/59,906)	(1,4	126/9,343)	(3,3	30/21,170)	(1,5	90/17,618)	(1,12	29/10,517)

^{*}Significant at the .05 level, two-sided test.

¹Models were estimated in a backward recurrence discrete-time survival framework with person-year the unit of analysis to predict the recency of most recent episode of the outcome disorders. Each model controlled for person-year, age-of-onset, time-since-onset, sex, race, and 19 dummy variables for the outcome disorder category.

2Model included all childhood SES variables and controlled for person-year, age category, time since

onset, sex, race, and diagnosis category.

³Model included covariates for the 11 CAs.

 $^{^4}$ n₁=the number of disorders included in the analysis; n_2 =the number of person-years included in the analysis.