
Appendix Table 1. Twelve-month treatment among respondents with Adult ADHD

	%	(se)
Specialty	25.9	4.4
General medical	18.8	3.4
Human services	7.4	1.5
CAM	18.4	5.1
Any treatment	42.4	4.4

¹CAM=complementary and alternative medicine

Appendix Table 2. WHO Disability Assessment Schedule (WHO-DAS) measures of 30-day disability associated with Adult ADHD (n=3199)¹

	% ²	(se) ²	OR ³	(95% CI) ³
I. Basic Functioning				
Self care	7.6	(2.7)	2.2*	(1.0-4.8)
Mobility	26.7	(4.7)	3.9*	(2.3-6.8)
Cognition	29.9	(5.3)	2.6*	(1.5-4.5)
II. Instrumental Functioning				
Time out of role	38.3	(4.9)	2.7*	(1.8-4.1)
Productive role functioning	35.0	(4.6)	2.1*	(1.4-3.2)
Social role functioning	18.6	(3.5)	3.5*	(2.1-5.9)

* Significant at the .05 level, two-sided design-based MI tests.

¹ Part II respondents ages 18-44

² Percent (standard error) of adults with ADHD who have the impairment

³ Based on bivariate logistic regression analysis using MI to estimate odds-ratios (OR) and 95% Confidence Intervals (95% CI)