

Table 1. Lifetime prevalence of DSM-IV/WMH-CIDI disorders by sex and cohort¹ (n=9282)

Lifetime	Total		Sex				Cohort							
			Female		Male		18-29		30-44		45-59		60+	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
I. Anxiety Disorders														
Panic disorder	4.7	(0.2)	6.2	(0.3)	3.1	(0.3)	4.2	(0.5)	5.9	(0.6)	5.9	(0.4)	2.1	(0.4)
Agoraphobia without panic	1.3	(0.1)	1.6	(0.2)	1.1	(0.2)	1.2	(0.3)	1.4	(0.2)	1.8	(0.3)	0.9	(0.2)
Specific phobia	12.5	(0.4)	15.8	(0.6)	8.9	(0.6)	13.0	(0.9)	13.9	(0.7)	14.4	(1.0)	7.7	(0.6)
Social phobia	12.1	(0.4)	13.0	(0.6)	11.1	(0.6)	13.3	(0.7)	14.5	(0.9)	12.6	(0.9)	6.8	(0.5)
Generalized anxiety disorder	5.7	(0.3)	7.1	(0.3)	4.2	(0.4)	4.3	(0.4)	6.5	(0.5)	7.6	(0.7)	4.0	(0.4)
Post-traumatic stress disorder ²	6.8	(0.4)	9.7	(0.7)	3.6	(0.3)	6.3	(0.6)	8.1	(0.9)	9.2	(0.8)	2.8	(0.5)
Obsessive-compulsive disorder ³	2.3	(0.3)	3.1	(0.5)	1.6	(0.3)	3.1	(0.7)	3.0	(0.9)	2.4	(0.8)	0.6	(0.3)
Adult/Child separation anxiety disorder ²	9.2	(0.4)	10.8	(0.6)	7.4	(0.5)	12.4	(0.9)	11.1	(0.7)	9.2	(0.8)	3.1	(0.5)
Any anxiety disorder ⁵	31.2	(1.0)	36.4	(1.1)	25.4	(1.2)	32.9	(1.3)	37.0	(1.5)	34.2	(1.7)	17.8	(1.4)
II. Mood Disorders														
Major depressive disorder	16.9	(0.5)	20.2	(0.5)	13.2	(0.8)	16.0	(0.8)	19.3	(0.9)	20.1	(1.2)	10.7	(0.7)
Dysthymia	2.5	(0.2)	3.1	(0.3)	1.8	(0.2)	1.8	(0.3)	2.8	(0.4)	3.8	(0.6)	1.3	(0.2)
Bipolar I-II-sub disorders	4.4	(0.3)	4.5	(0.3)	4.3	(0.4)	7.0	(0.8)	5.3	(0.4)	3.7	(0.4)	1.3	(0.3)
Any mood disorder	21.4	(0.6)	24.9	(0.6)	17.5	(0.9)	22.6	(1.0)	24.5	(1.0)	24.2	(1.2)	12.2	(0.9)
III. Impulse-control Disorders														
Oppositional-defiant disorder ⁴	8.5	(0.7)	7.7	(0.9)	9.3	(0.8)	9.9	(1.0)	7.3	(0.8)	--	--	--	--
Conduct disorder ⁴	9.5	(0.8)	7.1	(0.9)	12.0	(1.0)	10.8	(1.1)	8.4	(0.7)	--	--	--	--
Attention-deficit/hyperactivity disorder ⁴	8.1	(0.6)	6.4	(0.7)	9.8	(1.0)	7.8	(0.8)	8.3	(0.8)	--	--	--	--
Intermittent explosive disorder	7.4	(0.4)	5.7	(0.4)	9.2	(0.6)	12.6	(1.1)	8.8	(0.7)	5.3	(0.5)	2.4	(0.5)
Any impulse control disorder ⁴	25.0	(1.1)	21.6	(1.4)	28.6	(1.5)	27.0	(1.6)	23.4	(1.1)	--	--	--	--
IV. Substance Disorders														
Alcohol abuse with/without dependence ²	13.2	(0.6)	7.5	(0.5)	19.6	(0.9)	14.5	(1.0)	16.4	(1.1)	14.1	(1.0)	6.3	(0.7)
Drug abuse with/without dependence ²	8.0	(0.4)	4.8	(0.4)	11.6	(0.7)	11.1	(0.9)	12.1	(1.0)	6.8	(0.7)	0.3	(0.1)
Nicotine dependence ²	29.6	(0.8)	26.5	(1.3)	33.0	(1.0)	26.5	(1.8)	29.4	(1.5)	34.3	(1.6)	27.3	(1.7)
Any substance disorder ²	35.3	(0.9)	29.6	(1.3)	41.8	(1.1)	33.2	(1.9)	37.1	(1.8)	39.8	(1.5)	29.6	(1.7)
V. Any Disorder														
Any ⁵	57.4	(1.1)	56.5	(1.5)	58.4	(1.4)	58.7	(2.2)	63.7	(1.9)	60.0	(1.6)	44.0	(2.3)

¹This table includes updated data as of July 19, 2007. Updates reflect the latest diagnostic, demographic and raw variable information.

²Assessed in the Part II sample (n = 5692).

³Assessed in a random one-third of the Part II sample (n = 2073).

⁴Assessed in the Part II sample among respondents in the age range 18-44 (n = 3197).

⁵Estimated in the Part II sample. No adjustment is made for the fact that one or more disorders in the category were not assessed for all Part II respondents.