

09-24-2018

Auerbach, R. P., Mortier, P., Bruffaerts, R., Alonso, J., Benjet, C., Cuijpers, P., Demyttenaere, K., Ebert, D. D., Green, J. G., Hasking, P., Murray, E., Nock, M. K., Pinder-Amaker, S., Sampson, N. A., Stein, D. J., Vilagut, G., Zaslavsky, A. M., Kessler, R. C., & WHO WMH-ICS Collaborators. (2018). The WHO World Mental Health Surveys International College Student Project: Prevalence and Distribution of Mental Disorders. *Journal of Abnormal Psychology*. [Epub ahead of print].

The description of AUDIT scoring in the Methods section (under Measures/Mental Disorders) of this paper was incorrectly described as:

The version of the AUDIT we used, which defined alcohol use disorder as either a total score of 8+ or a score of 4+ on the AUDIT dependence questions (Babor, Higgins-Biddle, Saunders, & Monteiro, 2001), has been shown to have concordance with clinical diagnoses in the range AUC = 0.78-0.91 (Reinert & Allen, 2002).

This should read:

The version of the AUDIT we used, which defined alcohol use disorder as either a **total score of 16+ or a score 8-15 with 4+ on the AUDIT** dependence questions (Babor, Higgins-Biddle, Saunders, & Monteiro, 2001), has been shown to have concordance with clinical diagnoses in the range AUC = 0.78-0.91 (Reinert & Allen, 2002).

There was no error in the reported results.